



Unione
Matematica
Italiana



SBM
SOCIEDADE BRASILEIRA DE MATEMÁTICA



SIMAI
SOCIETÀ ITALIANA DI MATEMATICA
APPLICATA E INDUSTRIALE



SBMAC
Sociedade Brasileira de Matemática
Aplicada e Computacional



Università
degli Studi di
Messina

BOOK OF ABSTRACTS

SS10 - Fourier analysis and PDEs
on manifolds and Lie groups

2nd Joint Meeting Brazil-Italy
in Mathematics

September 9, 2026

Messina, Italy

Organizers:

Sandro Coriasco, University of Turin

Paulo L. Dattori da Silva, University of Sao Paulo at Sao Carlos
(ICMC-USP)

Alexandre Kirilov, Federal University of Paraná

Degenerate 3-evolution equations in Gevrey classes

Alexandre Arias Jr

Department of Computing and Mathematics, Universidade de São Paulo, Ribeirão Preto, Brazil

Alessia Ascanelli

Department of Mathematics and Computer Science, University of Ferrara, Italy

We consider the initial value problem for a 3–*evolution operator* P with variable coefficients and real characteristics, namely

$$(1) \quad P = D_t + a_3(t)D_x^3 + a_2(t, x)D_x^2 + a_1(t, x)D_x + a_0(t, x),$$

where $a_3 \in C([0, T], \mathbb{R})$, $a_j \in C([0, T]; \mathcal{B}^\infty(\mathbb{R}))$, $j = 0, 1, 2$, and $\mathcal{B}^\infty(\mathbb{R})$ is the space of complex-valued functions bounded on $\mathbb{R}$ together with all their derivatives.

We assume that $a_3(t)$ vanishes with finite order at $t = 0$:

$$\exists \ell > 0 : c_{a_3} t^\ell \leq |a_3(t)| \leq C_{a_3} t^\ell, \quad \forall t \in [0, T]$$

for some $0 < c_{a_3} \leq C_{a_3}$. In this situation, the equation degenerates and the (t, x) -behavior of the lower order terms becomes crucial to obtain well-posedness results. We assume the following Levi-type conditions: there exists positive constants $k, k', \sigma_1, \sigma_2$, such that for every $\beta \in \mathbb{N}$, $(t, x) \in [0, T] \times \mathbb{R}$

$$|\partial_x^\beta a_2(t, x)| \leq t^k C_{a_2}^{\beta+1} \beta! \langle x \rangle^{-\sigma_2}, \quad |\partial_x^\beta a_1(t, x)| \leq t^{k'} C_{a_1}^{\beta+1} \beta! \langle x \rangle^{-\sigma_1}, \quad |\partial_x^\beta a_0(t, x)| \leq C_{a_0}^{\beta+1} \beta!.$$

We show that:

- if a_2 degenerates in time with degeneracy gap $\ell - k > 0$, then the Cauchy problem is well-posed in suitable Gevrey-Sobolev spaces, and the (t, x) -behaviors of the coefficients (i.e., the parameters $k, k', \sigma_2, \sigma_1$) play together to determine well-posedness spaces;
- if only a_1 degenerates in time with degeneracy gap $\ell - k' > 0$, then Sobolev well-posedness can be recaptured if its decay as $x \rightarrow \infty$ is not-too-slow and $\ell - k'$ is small enough; otherwise, Gevrey-Sobolev well-posedness holds in spaces depending on $\sigma_2, \sigma_1, \ell, k'$;
- if there is no degeneracy in time ($k, k' \geq \ell$), only the x -decay rates σ_2, σ_1 determine the well posedness spaces, and Sobolev well-posedness can be recaptured if the two decays are not-too-slow.

References

- [1] A. Arias Jr, A. Ascanelli: *Degenerate 3-evolution equations in Gevrey classes*, submitted (2026), arXiv:2605.16576

Time-Periodic Evolution Equations

*Fernando de Ávila Silva*¹

Department of Mathematics, Federal University of Paraná

We discuss global properties of certain classes of evolution equations that are periodic in the time variable. The key tool in our approach is a characterization of the functional spaces in terms of Fourier analysis via eigenfunction expansions generated by a fixed elliptic differential operator. Hence, by employing a corresponding discretization process, we provide both necessary and sufficient conditions for the global solvability and global hypoellipticity of the equations under investigation.

The results are based on joint works with Alexandre Kirilov (UFPR-Brazil), Marco Cappiello (Unito-Italy), Matteo Bonino (Unito-Italy), and Sandro Coriasco (Unito-Italy).

References

- [1] F. de Ávila Silva, M. Bonino, S. Coriasco, *Global hypoellipticity and solvability for a class of evolution operators in time-periodic weighted Sobolev spaces*, Math. Ann, v. 394, (2026), p. 87.
- [2] F. de Ávila Silva, M. Cappiello, A. Kirilov, A. Global hypoellipticity for systems in time-periodic Gelfand-Shilov spaces, J. Funct. Anal., (2025), v. 290, p. 111300.
- [3] F. de Ávila Silva, M. Cappiello, *Globally solvable time-periodic evolution equations in Gelfand-Shilov classes*, Math. Ann, (2024), v. 391, p. 399-430.
- [4] F. de Ávila Silva, M. Cappiello, *Time-periodic Gelfand-Shilov spaces and global hypoellipticity on $\mathbb{T} \times \mathbb{R}^n$* , J. Funct. Anal. n. 9 (2022), 109418.
- [5] F. de Ávila Silva, T. V. Gramchev, A. Kirilov, *Global Hypoellipticity for First-Order Operators on Closed Smooth Manifolds*, J. Anal. Math., **135** (2018), 527–573.

¹The author thanks CNPq-Brazil and SBM-Brazil for their support.
E-mail: fernando.avila@ufpr.br.

Global solvability in time-periodic Gelfand-Shilov spaces for a class of first order operators

Adalberto P. Bergamasco

Departamento de Matemática, Instituto de Ciências Matemáticas e de Computação,
Universidade de São Paulo, São Carlos, Brazil

*Marco Cappiello*¹

Department of Mathematics, University of Turin, Italy

Nguyen T.H. Yen

Departamento de Matemática, Instituto de Ciências Matemáticas e de Computação,
Universidade de São Paulo, São Carlos, Brazil

Sérgio L. Zani

Departamento de Matemática, Instituto de Ciências Matemáticas e de Computação,
Universidade de São Paulo, São Carlos, Brazil

We investigate the global solvability in time-periodic Gelfand–Shilov spaces, cf. [1], for the first-order operator $L = \partial_t + a(x)\partial_x$ on $\mathbb{T} \times \mathbb{R}$, where $a(x)$ is a smooth real-valued function satisfying global Gevrey estimates on $\mathbb{R}$ and a suitable *slow decay* condition at infinity. We establish a complete characterization of global solvability in terms of the structure of the zero set of $a(x)$ in the non-compact setting $\mathbb{T} \times \mathbb{R}$. The same operator had been studied in [2] for $(t, x) \in \mathbb{T}^2$ and $a \in C^\infty(\mathbb{T}; \mathbb{R})$ in the C^∞ framework.

References

- [1] F. de Ávila Silva, M. Cappiello, *Time-periodic Gelfand-Shilov spaces and global hypoellipticity on $\mathbb{T} \times \mathbb{R}^n$* , J. Funct. Anal. **282** (2022), no. 9, 29 pp.
- [2] A.P. Bergamasco, G. Petronilho, *Closedness of the range for vector fields on the torus*, J. Differential Equations **154** (1999), 132–139.

¹Aknowledgements...

E-mail: marco.cappiello@unito.it.

Involutive systems on non-compact manifolds

*Sandro Coriasco*¹

Department of Mathematics “G. Peano”, University of Turin

Alexandre Kirilov

Department of Mathematics, Federal University of Paraná

Wagner Augusto Almeida de Moraes

Department of Mathematics, Federal University of Paraná

Pedro Meyer Tokoro

Department of Mathematics, Federal University of Paraná

We study the operator

$$(1) \quad L = dt + \sum_{k=1}^m \omega_k \wedge \partial_{x_k},$$

acting on differential forms over a product manifold of the form $M \times \mathbb{T}^m$, where M a smooth manifold and $\omega_1, \dots, \omega_m$ are smooth closed 1-forms on M . The characterisations of the hypoellipticity and solvability properties of operators of the form (1) are well-known when M is a closed manifold (that is, compact and without boundary; see, e.g., [1,2]). On the other hand, the cases where M is a non-compact manifold are not equally investigated. We show that when M is non-compact with a certain “structure at infinity”, it is still possible to characterise the global hypoellipticity and global solvability of L in terms of arithmetic properties of the forms $\omega_1, \dots, \omega_m$.

References

- [1] G. Araújo, P.L. Dattori da Silva, B. de Lessa Victor, *Global analytic hypoellipticity of involutive systems on compact manifolds*, Math. Ann. 386(3–4), (2023), 1321–1350.
- [2] A.P. Bergamasco, P.D. Cordaro, P.A. Malagutti, *Globally hypoelliptic systems of vector fields*, J. Funct. Anal. 114(2), (1993), 267–285.
- [3] S. Coriasco, A. Kirilov, W.A.A. de Moraes, P.M. Tokoro, *Global hypoellipticity for involutive systems on non-compact manifolds*, J. Geom. Anal. 36(22), (2026).

¹This work was supported in part by the Italian Ministry of the University and Research – MUR, within the framework of the Call relating to the scrolling of the final rankings of the PRIN 2022 – Project Code 2022HCLAZ8, CUP D53C24003370006 (PI A. Palmieri, Local unit Sc. Resp. S. Coriasco).

E-mail: sandro.coriasco@unito.it.

L^1 estimates for a dissipative system of elastic waves

Marcello D'Abbicco¹

Department of Mathematics, University of Bari

Guilherme de Martini

Department of Mathematics, Federal University of Santa Catarina

In this talk, we discuss some recent results concerning $L^p - L^q$ estimates for a dissipative system of elastic waves ($\mu > \lambda > 0$ and $\varepsilon > 0$):

$$(1) \quad \begin{cases} (\partial_{tt} - \lambda \Delta - (\mu - \lambda) \nabla \operatorname{div} + 2\varepsilon \partial_t) U = 0, & (t, x) \in (0, \infty) \times \mathbb{R}^n, \\ (U, U_t)(0, x) = (0, U_1)(x), & x \in \mathbb{R}^n, \end{cases}$$

with a special emphasis on L^1 estimates. A crucial role is played by the real Hardy space $H^1(\mathbb{R}^n)$, equipped with norm

$$\|f\|_{H^1} = \|f\|_{L^1} + \sum_{j=1}^n \|R_j f\|_{L^1},$$

where R_j are the Riesz transforms. In particular, the following case is of interest.

Theorem 1. *In space dimension $n = 2$, the following estimates hold:*

$$\begin{aligned} \|U(t, \cdot)\|_{L^1} &\leq C t (1 + \varepsilon t)^{-1} \|U_1\|_{L^1}, \\ \|U(t, \cdot)\|_{L^2} &\leq C \lambda^{-\frac{1}{2}} (\varepsilon t (1 + \varepsilon t))^{-\frac{3}{2}} \|U_1\|_{L^1} + e^{-\varepsilon t} \|U_1\|_{H^1}, \end{aligned}$$

where $C > 0$ independent on U_1 . In space dimension $n = 3$, the following estimate holds:

$$\|U(t, \cdot)\|_{L^1} \leq C t ((1 + \varepsilon t)^{-1} \|U_1\|_{L^1} + e^{-\varepsilon t} (1 + \varepsilon t)^{\frac{3}{2}} \|U_1\|_{H^1}).$$

¹The results have been partially obtained during the stay of the second author at the Department of Mathematics of University of Bari in the period September 2025-February 2026.

E-mail: marcello.dabbicco@uniba.it.

Global properties of tube-type systems of vector fields on compact manifolds

*Paulo L. Dattori da Silva*¹

University of Sao Paulo at Sao Carlos (ICMC-USP)

In this talk we will describe global properties of involutive structures defined on a smooth product manifold $M \times \mathbb{T}^m$, where $\mathbb{T}^m \simeq \mathbb{R}^n/2\pi\mathbb{Z}^m$ is the m -dimensional torus.

Given a compact n -dimensional manifold M and closed 1-forms $\omega_1, \dots, \omega_m$ over M , consider the product manifold $\Omega = M \times \mathbb{T}^m$ and the subbundle $\mathcal{V} \subset \mathbb{C}T\Omega$ annihilated by all the forms

$$\zeta_k = dx_k - \omega_k, \quad k \in \{1, \dots, m\},$$

where $(x_1, \dots, x_m)$ are angular coordinates on $\mathbb{T}^m$. Such a bundle is involutive, and gives rise to a complex of vector bundles over Ω and first-order differential operators between them.

Let Λ^1 be the bundle of complex 1-forms over M , and $\Lambda^{0,1}$ be its pullback to Ω via the natural projection $\Omega \rightarrow M$; we denote their spaces of smooth sections by $\Lambda^1 C^\infty(M)$ and $\Lambda^{0,1} C^\infty(\Omega)$, respectively. We define $\mathbb{L} : C^\infty(\Omega) \rightarrow \Lambda^{0,1} C^\infty(\Omega)$ by the formula

$$\mathbb{L}f = d_t f + \sum_{k=1}^m \omega_k \wedge \partial_{x_k} f,$$

where $t \in M$ and d_t is the exterior derivative on M ; hence, $\mathbb{L}$ is a first-order differential operator.

We will investigate the existence and regularity of global solutions of the equation $\mathbb{L}u = f$.

Also, we will consider related problems on the product $M \times G$, where G is a compact Lie group.

References

- [1] G. Araújo, P. L. Dattori da Silva, and B. de Lessa Victor, Global Analytic Solvability of Involutive Systems on Compact Manifolds. *Journal of Geometric Analysis*, v. 33, p. 151, 2023.
- [2] G. Araújo, P. L. Dattori da Silva, and B. de Lessa Victor, Global analytic hypoellipticity of involutive systems on compact manifolds. *Mathematische Annalen*, 2022.
- [3] P. L. Dattori da Silva and A. Meziani, A Gevrey differential complex on the torus. *Journal of Fourier Analysis and Applications*, 269(1): Paper No. 8, 2020.
- [4] P. L. Dattori da Silva, and F. M. Simão, A differential complex on a compact lie group. *Annali di Matematica* (2026). <https://doi.org/10.1007/s10231-026-01659-3>.
- [5] P. L. Dattori da Silva, F. M. Simão, and M. Ruzhansky, Global properties of involutive systems on compact manifolds, preprint.
- [6] I. V. Coelho da Silva, P. L. Dattori da Silva, and F. M. Simão, Regularity of solutions for a class of involutive systems of vector fields, preprint.

¹Supported in part by São Paulo Research Foundation (FAPESP), Brazil, Process Numbers #2024/08416-6 and #2024/12753-8 and National Council for Scientific and Technological Development, CNPq, Brazil (grant #313581/2021-5)

E-mail: dattori@icmc.usp.br.

Smoothing properties of third order operators with variable coefficients

Serena Federico

Department of Mathematics, University of Bologna

In this talk, we will discuss the smoothing properties of some third-order equations with variable coefficients modeling KdV-type equations in arbitrary space dimensions. We will describe the role of the nontrapping condition in the smoothing effect and relate it to our setting. Finally, well-posedness results for nonlinear initial value problems associated with this class of operators will be presented.

This talk is based on a joint work with D. Tramontana.

References

- [1] S. Federico, D. Tramontana, *Smoothing effect for third order operators with variable coefficients*. Preprint (2025). Arxiv. <https://arxiv.org/abs/2503.08656>

Semilinear Damped Wave Equations with Potential on Compact Lie Groups

Sandra Lucente

Department of Physics, University of Bari

In a recent project we study semilinear damped wave equations on compact Lie groups. More precisely, we consider a time-dependent potential in front of the power nonlinearity. The interaction between this potential and both the linear and nonlinear parts of the equation is exploited. In particular we discuss global existence result and lifespan estimates for blowing-up solution in the complementary case.

Global hypoellipticity of involutive systems on non-compact manifolds

Sandro Coriasco

Department of Mathematics, University of Turin

Alexandre Kirilov

Department of Mathematics, Federal University of Paraná

Wagner Augusto Almeida de Moraes

Department of Mathematics, Federal University of Paraná

*Pedro Meyer Tokoro*¹

Department of Mathematics, Federal University of Paraná

We study the global hypoellipticity of the operator

$$\mathbb{L} = d_t + \sum_{k=1}^m \omega_k \wedge \partial_{x_k},$$

defined on differential forms over a product manifold of the form $M \times \mathbb{T}^m$, where M is a non-compact manifold satisfying certain geometric and topological conditions, and $\omega_1, \dots, \omega_m$ are smooth closed 1-forms on M . Such operators are naturally associated with involutive subbundles of the complexified tangent bundle of $M \times \mathbb{T}^m$. Extending previous results obtained in the compact setting (see [1,2]), we characterize the global hypoellipticity and global solvability of $\mathbb{L}$ in terms of arithmetic properties of the forms $\omega_1, \dots, \omega_m$.

References

- [1] G. Araújo, P. L. Dattori da Silva, B. de Lessa Victor *Global analytic hypoellipticity of involutive systems on compact manifolds*, Math. Ann. 386(3–4), (2023), 1321–1350.
- [2] A. P. Bergamasco, P. D. Cordaro, P. A. Malagutti *Globally hypoelliptic systems of vector fields*, J. Funct. Anal. 114(2), (1993), 267–285.
- [3] S. Coriasco, A. Kirilov, W. A. A. de Moraes, P. M. Tokoro *Global hypoellipticity for involutive systems on non-compact manifolds*, J. Geom. Anal. 36(22), (2026).

¹This study was financed in part by CAPES – Brasil (Finance Code 001) and CNPq – Brasil (grant 420814/2025-6).

E-mail: pedro.tokoro@ufpr.br.

Global Properties of Operators on Compact Lie Groups and Beyond

Wagner Augusto Almeida de Moraes¹

Department of Mathematics, Federal University of Paraná

In this talk, we will discuss the global solvability and global hypoellipticity of left-invariant operators on compact Lie groups. Due to the fact that these operators act as Fourier multipliers, we can use the Fourier analysis based on irreducible unitary representations to study their properties through their symbols. We will start with a short historical background, move on to our recent contributions involving evolution equations and Vekua-type operators in different functional spaces, and point out possible future directions in closed manifolds in general and also for non-compact manifolds. Throughout the talk, we will illustrate these concepts with examples on the torus and $SU(2)$.

References

- [1] S. Coriasco, A. Kirilov, W. A. A. de Moraes, P. M. Tokoro, *Global hypoellipticity for involutive systems on non-compact manifolds*, The Journal of Geometric Analysis, 36 (2026), 22
- [2] A. Kirilov, A. P. Kowacs, W. A. A. de Moraes, *Global Solvability and Hypoellipticity for Evolution Operators on Tori and Spheres*, Mathematische Nachrichten, 297 (2024), 4605–4650
- [3] A. Kirilov, W. A. A. de Moraes, M. Ruzhansky, *Global Hypoellipticity and Global Solvability for Vector Fields on Compact Lie Groups*, Journal of Functional Analysis, 280 (2021), 108806
- [4] A. Kirilov, W. A. A. de Moraes, P. M. Tokoro, *Global solutions for systems of strongly invariant operators on closed manifolds*, Proceedings of the American Mathematical Society, 154 (2026), 1295–1310

¹The author was supported in part by CNPq-Brasil (Grant Nos. 420814/2025-6 and 301573/2025-5) and by Sociedade Brasileira de Matemática (SBM).

E-mail: wagnermoraes@ufpr.br.